

- [http://dx.doi.org/10.1016/S0021-9797\(03\)00393-X](http://dx.doi.org/10.1016/S0021-9797(03)00393-X)
- [5] P.C. Singer, 'DPBs in drinking water: additional scientific and policy considerations for public health protection' *J. Am. Water Works Assess.*, vol 98 (10), pp. 73-79, 2006.
- [6] X. Tan, X. Wang, C. Chen, and Sun, A., 'Effect of soil and fulvic acids, pH and ionic strength on Th (IV) sorption to TiO₂ nanoparticles,' *Applied Radiation and Isotopes*, vol 65, pp. 375-381, 2007. <http://dx.doi.org/10.1016/j.apradiso.2006.10.014>
- [7] J. Z. Xu, Q. H. Fan, Z. W. Niu, Y. Li, P. Li, and W. S. Wu, 'Studies of Eu(III) sorption on TiO₂ Effects of pH, humic acid and poly(acrylic acid),' *Chemical Engineering Journal*, vol 179, pp. 186-192, 2012. <http://dx.doi.org/10.1016/j.cej.2011.10.079>
- [8] H. Wang and J. P. Lewis, 'Second-generation photocatalytic materials anion-doped TiO₂,' *J. Phys: Condens. Matter*, vol 18, pp. 421-434, 2006. <http://dx.doi.org/10.1088/0953-8984/18/2/006>
- [9] P. G. Coble, S. A. Green, N. V. Blough, and R. B. Gagosian, 'Characterization of dissolved organic matter in the Black Sea by fluorescence spectroscopy,' *Nature*, vol 348 (6300), pp. 432-5, 1990. <http://dx.doi.org/10.1038/348432a0>
- [10] A. Baker 'Fluorescence excitation-emission matrix characterization of some sewage-impacted rivers,' *Environ Sci Technol*, vol 35(5) pp. 948-53, 2001. <http://dx.doi.org/10.1021/es000177t>
- [11] P. G. Coble, 'Characterization of marine and terrestrial DOM in seawater using excitation-emission matrix spectroscopy,' *Marine Chemistry*, vol 51, pp. 325-346, 1996. [http://dx.doi.org/10.1016/0304-4203\(95\)00062-3](http://dx.doi.org/10.1016/0304-4203(95)00062-3)
- [12] P. Akan, N.C. Birben, and M. Bekbolet, 'Adsorption of molecular size fractions of humic acid onto anion-doped TiO₂ Specimens,' *Desalination and Water Treatment*, vol 57(6), pp. 2455-2465, 2016. <http://dx.doi.org/10.1080/19443994.2015.1027277>
- [13] S. Kavurmaci Sen, and M. Bekbolet, 'Tracing TiO₂ photocatalytic degradation of humic acid in the presence of clay particles by excitation-emission matrix (EEM) fluorescence spectra' *Journal of Photochemistry and Photobiology A: Chemistry*, vol 282, pp. 53-61, 2014. <http://dx.doi.org/10.1016/j.jphotochem.2014.03.011>
- [14] P. Akan, N.C. Birben, and M., Bekbolet, 'Characterization of humic acid and its size fractions following adsorption onto anion-doped TiO₂ specimens by fluorescence spectroscopic studies,' *Proceedings of the World Congress on Civil, Structural, and Environmental Engineering (CSEE'16)*, Paper No: AWSPT129, International AET Inc, ISBN:978-1-927877-18-0, Prague, Czech Republic, March 30-31, 2016.
- [15] D.M. McKnight, E. W. Boyer, P.K. Westerhoff, P.T. Doran, T. Kulbe, and D.T. Anderson, 'Spectrofluorometric characterization of dissolved organic matter for indication of precursor organic material and aromaticity,' *Limnol. Oceanogr.*, vol 46, pp. 38-48, 2001. <http://dx.doi.org/10.4319/lo.2001.46.1.0038>
- [16] T. F. Marhaba, K. Bengraïne, Y. Pu, and J. Arago, 'Spectral fluorescence signatures and partial least squares regression: model to predict dissolved organic carbon in water,' *J. Hazard. Mater*, vol 97, pp. 83-97, 2003. [http://dx.doi.org/10.1016/S0304-3894\(02\)00246-7](http://dx.doi.org/10.1016/S0304-3894(02)00246-7)



Perihan AKAN: I am a research assistant and also PhD. student at Department of Environmental Engineering, Hacettepe University, Ankara/Turkey. I took my master degree from Bogazici University, Institute of Environmental Sciences, Istanbul/Turkey. I graduated from Environmental Engineering Department, Trakya University, Edirne/Turkey. My research areas are advanced oxidation processes, adsorption as well as air pollution control.

ACKNOWLEDGEMENT

Financial support provided by Research Fund of Bogazici University through Project No: 6750 is gratefully acknowledged.